

The induced representation of the isometry group of the Euclidean Taub-NUT space and new spherical harmonics

Ion I. Cotăescu *

*West University of Timișoara,
V. Pârvan Ave. 4, RO-1900 Timișoara, Romania*

Mihai Visinescu †

*Department of Theoretical Physics,
National Institute for Physics and Nuclear Engineering,
P.O.Box M.G.-6, Magurele, Bucharest, Romania*

February 1, 2008

Abstract

It is shown that the $SO(3)$ isometries of the Euclidean Taub-NUT space combine a linear three-dimensional representation with one induced by a $SO(2)$ subgroup, giving the transformation law of the fourth coordinate under rotations. This explains the special form of the angular momentum operator on this manifold which leads to a new type of spherical harmonics and spinors.

Keywords: *Euclidean Taub-NUT space, spherical harmonics, spherical spinors, isometry group, induced representation.*

Pacs 04.62.+v

*E-mail: cota@physics.uvt.ro

†E-mail: mvisin@theor1.theory.nipne.ro

1 Introduction

The Euclidean Taub-Newman-Unti-Tamburino (Taub-NUT) metric is involved in many modern studies in physics. One of the first examples given of a gravitational instanton was the self-dual Taub-NUT solution [1]. Much attention has been paid to the Euclidean Taub-NUT metric since in the long-distance limit the relative motion of two monopoles is described approximately by its geodesics [2, 3]. On the other hand this metric is just the space part of the metric of the celebrated Kaluza-Klein monopole of Gross and Perry [4] and of Sorkin [5].

The Taub-NUT space is also of interest since beside isometries there are hidden symmetries giving rise to conserved quantities associated to Stäckel-Killing tensors [6]. There is a conserved vector, analogous to the Runge-Lenz vector of the Kepler type problem, whose existence is rather surprising in view of the complexity of the equations of motion [7]-[10]. These hidden symmetries are related to the existence of special objects arising in this geometry, i.e. four Killing-Yano tensors [11] generating the Stäckel-Killing ones [8], [12]-[14].

The quantum theory in the Euclidean Taub-NUT background has also interesting specific features in the case of the scalar fields [15] as well as for Dirac fields of spin $\frac{1}{2}$ fermions [16]-[18]. In both cases there exist large algebras of conserved observables [19] including the components of the angular momentum and three components of the Runge-Lenz operator that complete a six-dimensional dynamical algebra [6, 18, 19]. Remarkably, the orbital angular momentum has a special unusual form that generates new harmonics called $SO(3) \otimes U(1)$ -harmonics [15]. What is the explanation of this fact?

In our opinion the form of the angular momentum operator is determined by the specific type of isometries of the Euclidean Taub-NUT space which combine linear representations with induced ones [20, 21]. The purpose of this article is to prove this showing that the fourth Cartesian (or spherical) coordinate transforms under rotations according to a representation of the $SO(3)$ isometry group induced by one of its $SO(2)$ subgroups. Thus, after we present in the next section the form of the isometry transformations, we demonstrate in Sec. 3 that these involve an induced representation for which we deduce the explicit closed form of the transformation law. The Sec. 4 is devoted to the isometry generators showing how the induced representation determines the specific form of the angular momentum operators of the scalar or spin half theories. The corresponding new spherical harmonics [15] as well

as the associated Pauli spherical spinors [16] are briefly presented in the next section. Some technical details are given in an Appendix.

2 $SO(3) \otimes U(1)$ isometry transformations

The Euclidean Taub-NUT manifold M_4 is a 4-dimensional Kaluza-Klein space which has static charts with the Cartesian coordinates x^μ ($\mu, \nu, \dots = 1, 2, 3, 4$) where x^i ($i, j, \dots = 1, 2, 3$) are the *physical* Cartesian space coordinates while x^4 is the Cartesian extra-coordinate. In the usual three-dimensional vector notations, $\mathbf{x} = (x^1, x^2, x^3)$, $r = |\mathbf{x}|$ and $dl^2 = d\mathbf{x} \cdot d\mathbf{x}$, its line element reads

$$ds^2 = \frac{1}{V(r)} dl^2 + V(r) [dx^4 + A_i(\mathbf{x}) dx^i]^2 \quad (1)$$

where

$$\frac{1}{V} = 1 + \frac{\mu}{r}, \quad A_1 = -\frac{\mu}{r} \frac{x^2}{r + x^3}, \quad A_2 = \frac{\mu}{r} \frac{x^1}{r + x^3}, \quad A_3 = 0. \quad (2)$$

The real number μ is the main parameter of the theory. If one interprets, $\mathbf{A}$ as the gauge field of a monopole it results the magnetic field with central symmetry

$$\mathbf{B} = \mu \frac{\mathbf{x}}{r^3}. \quad (3)$$

Other important charts are those with spherical coordinates (r, θ, ϕ, χ) where r, θ, ϕ , are commonly related to the physical Cartesian ones, x^i . The apparent singularity at the origin is unphysical if x^4 is periodic with period $4\pi\mu$ [7]-[10]. Consequently the fourth coordinate χ is defined such that

$$x^4 = -\mu(\chi + \phi). \quad (4)$$

This chart covers the domain where $r > 0$ if $\mu > 0$ or $r > |\mu|$ if $\mu < 0$, the angular coordinates θ, ϕ cover the sphere S^2 and $\chi \in D_\chi = [0, 4\pi)$. The line element in spherical coordinates is

$$ds^2 = \frac{1}{V} (dr^2 + r^2 d\theta^2 + r^2 \sin^2 \theta d\phi^2) + \mu^2 V (d\chi + \cos \theta d\phi)^2, \quad (5)$$

since

$$A_r = A_\theta = 0, \quad A_\phi = \mu(1 - \cos \theta). \quad (6)$$

The Euclidean Taub-NUT spaces M_4 possess a special type of isometries which combines the space transformations with the gauge transformations of the gauge field $\mathbf{A}(\mathbf{x})$. There are $U_4(1)$ transformations $x^4 \rightarrow x'^4 = x^4 + \lambda$ which leave the metric invariant if λ is a point-independent real constant. Moreover, if one takes $\lambda = \lambda(\mathbf{x})$ an arbitrary function of $\mathbf{x}$ then these become gauge transformations preserving the form of the line element only if one requires $\mathbf{A}$ to transform as

$$A_i(\mathbf{x}) \rightarrow A'_i(\mathbf{x}) = A_i(\mathbf{x}) - \partial_i \lambda(\mathbf{x}). \quad (7)$$

Thus it is obvious that $U_4(1)$ is an isometry group playing, in addition, the role of the gauge group associated to the gauge field $\mathbf{A}$. In other respects, this geometry allows an $SO(3)$ symmetry given by usual *linear* rotations of the physical space coordinates, $\mathbf{x} \rightarrow \mathbf{x}' = R\mathbf{x}$ with $R \in SO(3)$, and the special non-linear transformations of the fourth coordinate,

$$R : \quad x^4 \rightarrow x'^4 = x^4 + h(R, \mathbf{x}), \quad (8)$$

produced by a function h depending on R and $\mathbf{x}$ which must satisfy

$$h(R'R, \mathbf{x}) = h(R', R\mathbf{x}) + h(R, \mathbf{x}), \quad h(Id, \mathbf{x}) = 0 \quad (9)$$

where Id is the identity of $SO(3)$. Obviously, this condition guarantees that Eq. (8) defines a representation of the $SO(3)$ group. These transformations preserve the general form of the line element (1) if $\mathbf{A}$ transforms *manifestly* covariant under rotations as a vector field, up to a gauge transformation, V being a scalar. In this way one obtains a representation of the group $SO(3) \otimes U_4(1)$ whose transformations,

$$\mathbf{x} \rightarrow \mathbf{x}' = R\mathbf{x} \quad (10)$$

$$[R, \lambda(\mathbf{x})] : \quad x^4 \rightarrow x'^4 = x^4 + h(R, \mathbf{x}) + \lambda(\mathbf{x}) \quad (11)$$

$$\mathbf{A}(\mathbf{x}) \rightarrow \mathbf{A}'(\mathbf{x}') = R \{ \mathbf{A}(\mathbf{x}) - \nabla [h(R, \mathbf{x}) + \lambda(\mathbf{x})] \}, \quad (12)$$

produced by any $R \in SO(3)$ and real function $\lambda(\mathbf{x})$, combines isometries and gauge transformations. Hereby we can *separate* the isometries requiring that for point-independent parameters λ the components of the gauge field remain unchanged, i.e. $A'_i = A_i$. According to Eq. (12), this condition can be written as

$$\nabla h(R, \mathbf{x}) = \mathbf{A}(\mathbf{x}) - R^{-1} \mathbf{A}(R\mathbf{x}) \quad (13)$$

defining the *specific* function h corresponding to the gauge field $\mathbf{A}$. Therefore, the isometry transformations $(R, \lambda) : x \rightarrow x'$ are three-dimensional rotations and x^4 translations that transform $x = (\mathbf{x}, x^4)$ into $x' = (\mathbf{x}', x'^4)$ according to Eqs. (10) and (11) restricted to point-independent values of λ and function h defined by Eq. (13). These form the isometry group $I(M_4) = SO(3) \otimes U_4(1)$ of the Euclidean Taub-NUT space M_4 . What is remarkable here is that the representation of $I(M_4)$ carried by M_4 mixes up linear transformations with non-linear ones involving the function h .

In our opinion the study of this type of representation is important since it governs the transformation laws of the vectors and tensors under isometries that are the starting points in deriving conserved quantities through the Noether theorem. For example, the isometry (R, λ) transforms the contravariant components $v^\mu(x)$ of a four-dimensional vector field as

$$v^i(x) \rightarrow v'^i(x') = R^i_j v^j(x), \quad (14)$$

$$v^4(x) \rightarrow v'^4(x') = v^4(x) + v^i(x) \partial_i h(R, \mathbf{x}). \quad (15)$$

It is worth pointing out that from Eqs. (13) and (15) we can deduce that the quantity

$$v'^4(x') + \mathbf{A}(\mathbf{x}') \cdot \mathbf{v}'(x') = v^4(x) + \mathbf{A}(\mathbf{x}) \cdot \mathbf{v}(x) \quad (16)$$

behaves as a scalar. These interesting properties of isometries could be better understood finding explicitly the analytical expression of the function h .

3 The induced representation

The study of the function h must combine the integration of the equations (13) with some algebraic properties resulted from the condition (9). The main point is to show that the transformation rule (8) of the fourth coordinate of M_4 is given by a representation of the isometry group induced by one of its subgroups. We recall that the $SO(3)$ subgroup of $I(M_4)$ has three independent one-parameter subgroups, $SO_i(2)$, $i = 1, 2, 3$, each one including rotations $R_i(\psi)$, of angles $\psi \in [0, 2\pi)$ around the axis i . With this notation any rotation $R \in SO(3)$ in the usual Euler parametrization reads $R(\alpha, \beta, \gamma) = R_3(\alpha)R_2(\beta)R_3(\gamma)$.

We start with the observation that because of the special form of gauge field (2) all the rotations of the subgroup $SO_3(2)$ satisfy the condition

$$\hat{R}\mathbf{A}(\mathbf{x}) = \mathbf{A}(\hat{R}\mathbf{x}), \quad \forall \hat{R} \in SO_3(2). \quad (17)$$

Bearing in mind that, in addition, the gauge field does not depend on x^4 we say that the subgroup $H(M_4) = SO_3(2) \otimes U_4(1) \subset I(M_4)$ is by definition the *little group* associated to $\mathbf{A}$. In what follows we are interested to exploit the existence of the little group focusing on the rotations $\hat{R} \in SO_3(2)$. According to Eqs. (13) and (17) it results that

$$h(\hat{R}, \mathbf{x}) \equiv \hat{h}(\hat{R}) \quad (18)$$

is point-independent being defined only on $SO_3(2)$. Then the condition (9) becomes

$$\hat{h}(\hat{R}\hat{R}') = \hat{h}(\hat{R}) + \hat{h}(\hat{R}'), \quad \forall \hat{R}, \hat{R}' \in SO_3(2) \quad (19)$$

which means that the set of values $\{\hat{h}(\hat{R}) \mid \hat{R} \in SO_3(2)\}$ form a one-dimensional representation of $SO_3(2)$ provided $\hat{h}(Id) = 0$. This representation is non-trivial (with $\hat{h}(\hat{R}) \neq 0$ when $\hat{R} \neq Id$) only if we assume that

$$\hat{h}[R_3(\alpha)] = \text{const.} \alpha. \quad (20)$$

These properties suggest us to try to find the function h using rotations in the Euler parametrization and the chart with spherical coordinates where the differential equations could be simpler since $h(R, \mathbf{x}) = h(R, \theta, \phi)$ does not depend on the radial coordinate r . Then, according to Eqs. (18) and (19), we can write

$$\begin{aligned} h[R(\alpha, \beta, \gamma), \mathbf{x}] &= \hat{h}[R_3(\alpha)] + h[R_2(\beta)R_3(\gamma), \mathbf{x}] \\ &= h[R_2(\beta), R_3(\gamma)\mathbf{x}] + \hat{h}[R_3(\alpha + \gamma)] \end{aligned} \quad (21)$$

pointing out that the central problem is to integrate the system (13) in spherical coordinates for the particular case of $R = R_2(\beta)$. Denoting $h[R_2(\beta), \mathbf{x}] \equiv h(\beta, \theta, \phi)$, after a few manipulation we find that Eqs. (13) take the form

$$\partial_\theta h(\beta, \theta, \phi) = -\mu \frac{\sin \phi \sin \beta}{1 + \cos \theta \cos \beta - \sin \theta \cos \phi \sin \beta}, \quad (22)$$

$$\partial_\phi h(\beta, \theta, \phi) = \mu \frac{(1 - \cos \theta)(1 - \cos \beta) - \sin \theta \cos \phi \sin \beta}{1 + \cos \theta \cos \beta - \sin \theta \cos \phi \sin \beta}. \quad (23)$$

The integration of this system gives $h(\beta, \theta, \psi)$ up to some arbitrary integration constants as it is shown in the Appendix. However, the definitive form of the function $h(R, \theta, \phi)$ can be easily found by using the technique of induced representations.

Let us consider the chart with spherical coordinates (r, θ, ϕ, χ) where θ and ϕ can be interpreted as the Euler angles of the rotation giving $\mathbf{x} = R(\theta, \phi, 0)\mathbf{x}_o$ from $\mathbf{x}_o = (0, 0, r)$. After an arbitrary rotation $R(\alpha, \beta, \gamma) \in I(M_4)$ we arrive to the chart with the new coordinates $(r, \theta', \phi', \chi')$ among them the first three are the spherical coordinates of the transformed vector

$$\begin{aligned}\mathbf{x}' &= R(\phi', \theta', 0)\mathbf{x}_o = R(\alpha, \beta, \gamma)\mathbf{x} \\ &= R(\alpha, \beta, \gamma)R(\phi, \theta, 0)\mathbf{x}_o.\end{aligned}\tag{24}$$

In addition, we assume that the fourth transformed spherical coordinate χ' is defined such that

$$R(\phi', \theta', \chi') = R(\alpha, \beta, \gamma)R(\phi, \theta, 0)\tag{25}$$

from which we deduce

$$\begin{aligned}R_3(\chi') &= R^{-1}(\phi', \theta', 0)R(\alpha, \beta, \gamma)R(\phi, \theta, 0) \\ &= R^{-1}(\phi' - \alpha, \theta', 0)R_2(\beta)R(\phi + \gamma, \theta, 0).\end{aligned}\tag{26}$$

Thus we obtain the transformation $R : \chi \rightarrow \chi'$ of a representation of the $SO(3)$ group *induced* by the subgroup $SO_3(2)$. The corresponding induced representation transforming the fourth Cartesian coordinate x^4 is defined by Eq. (8) where the function h must be

$$h(R, \theta, \phi) = -\mu(\chi' + \phi' - \phi).\tag{27}$$

The last step is to rewrite Eq. (26) in terms of $SU(2)$ transformations corresponding to all the particular rotations involved therein. In this way it is not difficult to find the final result:

$$\begin{aligned}h[R(\alpha, \beta, \gamma), \theta, \phi] &= -\mu(\alpha + \gamma) \\ &\quad - 2\mu \arctan \left[\frac{\sin(\phi + \gamma)}{\cot \frac{\theta}{2} \cot \frac{\beta}{2} - \cos(\phi + \gamma)} \right].\end{aligned}\tag{28}$$

Now we can convince ourselves that this result is correct since for $\alpha = \gamma = 0$ the function $h[R(0, \beta, 0), \theta, \phi] = h(\beta, \theta, \phi)$ is the desired solution of the system (22) - (23) calculated in Appendix. Moreover, if we put $\beta = \gamma = 0$ we obtain that the point-independent functions (20) are $h[R_3(\alpha), \theta, \phi] = \hat{h}[R_3(\alpha)] = -\mu \alpha$ (with $\text{const.} = -\mu$).

4 The angular momentum

In the quantum theory on the Euclidean Taub-NUT background the basic operators are introduced using the geometric quantization. Now when we know the closed form of the function h we can calculate the generators of the natural representation of the group $I(M_4)$ using only group theoretical methods.

The natural representation is carried by the space of scalar wave functions Φ defined on M_4 which have the transformation rule $(R, \lambda) : \Phi(x) \rightarrow \Phi'(x') = \Phi(x)$ under isometries $(R, \lambda) \in I(M_4)$. In this representation the generators are the (orbital) differential operators defined as usual. More precisely, denoting by ξ a set of parameters of $I(M_4)$ and by $x'(x, \xi)$ the transformed coordinates, one associates to each parameter ξ^a the Killing vector k_a of components

$$k_a^\mu(x) = \left. \frac{\partial x'^\mu(x, \xi)}{\partial \xi^a} \right|_{\xi=0}, \quad (29)$$

and the vector field

$$X_a = -ik_a^\mu(x) \partial_\mu \quad (30)$$

that is just the generator of the natural representation corresponding to ξ^a . In the case of our isometry group $I(M_4)$ we take the first three parameters, ξ^i , the Cayley-Klein parameters of the one-parameter subgroups $SO_i(2)$ and $\xi^4 = \lambda$. Then we find that the generator of the $U_4(1)$ translations is the fourth component of the momentum operator, $P_4 = -i\partial_4$. Furthermore, exploiting the form of h we calculate the rotation generators in Cartesian coordinates. These are the components of the orbital angular momentum operator,

$$L_1 = -i(x^2\partial_3 - x^3\partial_2) + i\mu \frac{x^1}{r + x^3} \partial_4 \quad (31)$$

$$L_2 = -i(x^3\partial_1 - x^1\partial_3) + i\mu \frac{x^2}{r + x^3} \partial_4 \quad (32)$$

$$L_3 = -i(x^1\partial_2 - x^2\partial_1) + i\mu \partial_4. \quad (33)$$

The last terms arising from Eqs. (A.5) are due to the existence of the function h giving the induced representation discussed above. Thus we recover the known expression of the angular momentum operator

$$\mathbf{L} = \mathbf{x} \times \mathbf{P} - \mu \frac{\mathbf{x}}{r} P_4 \quad (34)$$

where $\mathbf{P} = -i(\boldsymbol{\partial} - \mathbf{A}\partial_4)$ is the three-dimensional (physical) momentum whose components obey the commutation rules $[P_i, P_j] = i\epsilon_{ijk}B_kP_4$ and $[P_i, P_4] = 0$.

The scalar quantum mechanics in the Taub-NUT geometry [10] is based on the Schrödinger or Klein-Gordon equations involving the static operator

$$\Delta = -\nabla_\mu g^{\mu\nu} \nabla_\nu = V\mathbf{P}^2 + \frac{1}{V}P_5^2 \quad (35)$$

which is either proportional with the Hamiltonian operator of the Schrödinger theory ($\Delta = 2H$) or represents the static part of the Klein-Gordon operator [19]. In both cases when an operator commutes with Δ we say that this is conserved. One can verify that the operators L_i are conserved and satisfy the canonical commutation rules. Moreover, their commutators with the coordinates and the momentum operators are the usual ones.

Other three important conserved operators are given by the three specific Killing tensors of the Taub-NUT geometry, $\mathbf{k}_{\mu\nu}$ [10]. With their help one defines the vector operator which plays the same role as the Runge-Lenz vector in the usual quantum mechanical Kepler problem:

$$\mathbf{K} = -\frac{1}{2}\nabla_\mu \mathbf{k}^{\mu\nu} \nabla_\nu = \frac{1}{2}(\mathbf{P} \times \mathbf{L} - \mathbf{L} \times \mathbf{P}) - \mu \frac{\mathbf{x}}{r} \left(\frac{1}{2}\Delta - \frac{1}{\mu^2}Q^2 \right) \quad (36)$$

with

$$Q = -\mu P_4 = -i\partial_\chi. \quad (37)$$

This transforms as a vector under space rotations since its components obey

$$[L_i, K_j] = i\epsilon_{ijk}K_k. \quad (38)$$

Moreover, the operators K_i can be rescaled such that in association with the operators L_i should constitute the basis generators of different dynamical algebras, $so(4)$, $so(3,1)$ or $e(3)$, corresponding to different domains of energy of the free quantum motion of scalar particles in the Taub-NUT background [10]. In our opinion, all this machinery works because of the special form of the operators L_i whose supplemental terms are produced by the induced representation discussed here.

When one goes to the chart with spherical coordinate a special attention must be paid to the definition (4) which shows that the fourth spherical coordinate χ is *translated* with ϕ . This means that the whole algebra of observables in Cartesian coordinates must be transformed according to this

translation into an *equivalent* algebra. To do this it is convenient to define the translation operator in terms of the operator Q (37):

$$U(\phi) = e^{i\phi Q} \quad (39)$$

which has to transform any operator X defined in the Cartesian chart, giving the corresponding observable $\tilde{X} = U(\phi)XU^\dagger(\phi)$ in the spherical chart. In other words in the spherical charts we use a representation of the operator algebra different from that considered in Cartesian charts. Thus the components of the orbital angular momentum (34) in the canonical basis (with $\tilde{L}_\pm = \tilde{L}_1 \pm i\tilde{L}_2$) become

$$\tilde{L}_3 = -i\partial_\phi, \quad (40)$$

$$\tilde{L}_\pm = e^{\pm i\phi} \left[\pm \partial_\theta + i \left(\cot \theta \partial_\phi - \frac{1}{\sin \theta} \partial_\chi \right) \right]. \quad (41)$$

Many other operators including the Runge-Lenz vector will take new forms in this representation but preserving their commutation relations.

In theories with spin the form of the conserved operators associated with isometries is, in addition, strongly dependent on the gauge one uses [22, 23]. The space M_4 allows a special Cartesian gauge in the chart with Cartesian coordinates [24] which leads to a total angular momentum having standard components $J_i = L_i + S_i$, with point-independent spin operators S_i [16]. The corresponding operators in the spherical chart are $\tilde{J}_i = \tilde{L}_i + S_i$ since S_i commute with $U(\phi)$. We note that the Runge-Lenz operator in theories with spin half has specific spin terms as we have recently pointed out [17, 18]. It is interesting that the basic commutation relations of the algebra of observables are the same in the scalar theory as well as in theories with spin $\frac{1}{2}$ of Pauli or Dirac type [18].

5 New harmonics and spherical spinors

It is natural that the unusual form of the $SO(3)$ generators (40) and (41) should lead to new spherical harmonics. These have to be the common eigenfunctions of complete set of commuting operators $\{\tilde{\mathbf{L}}^2, \tilde{L}_3, Q\}$ representing a basis of the Hilbert space of square integrable functions of ϕ , θ and χ defined on the compact domain $S^2 \times D_\chi$. In Ref. [15] we introduced the

$SO(3) \otimes U(1)$ -harmonics, $Y_{l,m}^q$, which satisfy the eigenvalue problems

$$\tilde{\mathbf{L}}^2 Y_{l,m}^q = l(l+1) Y_{l,m}^q, \quad (42)$$

$$\tilde{L}_3 Y_{l,m}^q = m Y_{l,m}^q, \quad (43)$$

$$Q Y_{l,m}^q = q Y_{l,m}^q, \quad (44)$$

and the orthonormalization condition

$$\begin{aligned} \langle Y_{l,m}^q, Y_{l',m'}^{q'} \rangle &= \int_{S^2} d(\cos \theta) d\phi \int_0^{4\pi} d\chi Y_{l,m}^q(\theta, \phi, \chi)^* Y_{l',m'}^{q'}(\theta, \phi, \chi) \\ &= \delta_{l,l'} \delta_{m,m'} \delta_{q,q'}, \end{aligned} \quad (45)$$

The boundary conditions on $S^2 \times D_\chi$ require l and m to be integer numbers while $q = 0, \pm 1/2, \pm 1, \dots$ [10] but, in general, q can be any real number. Solving Eqs. (43) and (44) we get

$$Y_{l,m}^q(\theta, \phi, \chi) = \frac{1}{4\pi} \Theta_{l,m}^q(\cos \theta) e^{im\phi} e^{iq\chi} \quad (46)$$

where the function $\Theta_{l,m}^q$ calculated according to Eq. (42) must satisfy the normalization condition

$$\int_{-1}^1 d(\cos \theta) \left| \Theta_{l,m}^q(\cos \theta) \right|^2 = 2. \quad (47)$$

resulted from Eq. (45). This problem has solutions for any values of the quantum numbers obeying

$$|q| - 1 < |m| \leq l \quad (48)$$

when one finds [15]

$$\begin{aligned} \Theta_{l,m}^q(\cos \theta) &= \frac{\sqrt{2l+1}}{2^{|m|}} \left[\frac{(l-|m|)!(l+|m|)!}{\Gamma(l-q+1)\Gamma(l+q+1)} \right]^{\frac{1}{2}} \\ &\times (1 - \cos \theta)^{\frac{|m|-q}{2}} (1 + \cos \theta)^{\frac{|m|+q}{2}} P_{l-|m|}^{(|m|-q, |m|+q)}(\cos \theta). \end{aligned} \quad (49)$$

For $m = |m|$ the $SO(3) \otimes U(1)$ harmonics are given by (46) and (49) while for $m < 0$ we have to use the obvious formula

$$Y_{l,-m}^q = (-1)^m (Y_{l,m}^{-q})^*. \quad (50)$$

When the boundary conditions allow half-integer quantum numbers l and m then we say that the functions defined by Eqs. (46) and (49) (up to a suitable factor) represent $SU(2) \otimes U(1)$ harmonics.

Thus we have obtained a non-trivial generalization of the spherical harmonics of the same kind as the spin-weighted spherical harmonics [25] or those studied in [26]. Indeed, if l , m and $q = m'$ are either integer or half-integer numbers then we have

$$Y_{l,m}^{m'}(\theta, \phi, \chi) = \frac{\sqrt{2l+1}}{4\pi} D_{m,m'}^l(\phi, \theta, \chi) \quad (51)$$

where $D_{m,m'}^l$ are the matrix elements of the irreducible representation of weight l of the $SU(2)$ group corresponding to the rotation of Euler angles (ϕ, θ, χ) . What is new here is that our harmonics are defined for any real number q . For this reason these are useful in solving some actual physical problems [27]. Notice that similar spherical harmonics were used recently in [28] under a different name (ring-shaped harmonics)¹.

With the help of these spherical harmonics we constructed the corresponding spherical spinors [16] following the traditional method [29]. We defined the spherical spinors $\Psi_{q,j,m_j}^\pm(\theta, \phi, \chi)$ as the common two-component eigenspinors of the eigenvalue problems

$$Q \Psi_{q,j,m_j}^\pm = q \Psi_{q,j,m_j}^\pm, \quad (52)$$

$$\tilde{\mathbf{J}}^2 \Psi_{q,j,m_j}^\pm = j(j+1) \Psi_{q,j,m_j}^\pm, \quad (53)$$

$$\tilde{J}_3 \Psi_{q,j,m_j}^\pm = m_j \Psi_{q,j,m_j}^\pm, \quad (54)$$

$$(\sigma_L + 1_{2 \times 2}) \Psi_{q,j,m_j}^\pm = \pm(j+1/2) \Psi_{q,j,m_j}^\pm, \quad (55)$$

with $\sigma_L = \sigma_i \tilde{L}_i$ where σ_i are the Pauli matrices while $1_{2 \times 2}$ is the 2×2 identity matrix. These spinors are, in addition, eigenfunctions of $\tilde{\mathbf{L}}^2$ corresponding to the eigenvalues $l(l+1)$ with $l = j \pm \frac{1}{2}$. For $j = l + \frac{1}{2} > |q| - \frac{1}{2}$ we have [29, 15]

$$\Psi_{q,j,m_j}^+ = \frac{1}{\sqrt{2j}} \begin{pmatrix} \sqrt{j+m_j} Y_{j-\frac{1}{2},m_j-\frac{1}{2}}^q \\ \sqrt{j-m_j} Y_{j-\frac{1}{2},m_j+\frac{1}{2}}^q \end{pmatrix} \quad (56)$$

¹The author seems to be not aware of the previous results of [15].

while for $j = l - \frac{1}{2} > |q| - \frac{3}{2}$ we get

$$\Psi_{q,j,m_j}^- = \frac{1}{\sqrt{2j+2}} \begin{pmatrix} \sqrt{j-m_j+1} Y_{j+\frac{1}{2},m_j-\frac{1}{2}}^q \\ -\sqrt{j+m_j+1} Y_{j+\frac{1}{2},m_j+\frac{1}{2}}^q \end{pmatrix}. \quad (57)$$

These spherical spinors are orthonormal since the $SO(3) \otimes U(1)$ harmonics are orthonormal with respect to the angular scalar product (45).

Finally, denoting $\sigma_r = \sigma_i x^i / r$ and using Eq.(55) and the relation

$$\{\sigma_r, \sigma_L + 1_{2 \times 2}\} = 2q 1_{2 \times 2} \quad (58)$$

we find that

$$\sigma_r \Psi_{q,j,m_j}^\pm = \pm \lambda_j^q \Psi_{q,j,m_j}^\pm + \sqrt{1 - (\lambda_j^q)^2} \Psi_{q,j,m_j}^\mp \quad (59)$$

where $\lambda_j^q = q/(j + \frac{1}{2})$. Note that a similar result is obtained in Ref. [30]. By using this property we have written down the particular solutions of the Dirac equation in the central charts of the Taub-NUT background [16].

6 Conclusions

We showed here how transforms under rotations the fourth coordinate of the Euclidean Taub-NUT space pointing out that the transformation law is of a representation of the group $SO(3)$ induced by one of its $SO(2)$ subgroups. In this context the angular momentum operators get new specific terms that do not change their algebraic relations but give rise to new $SO(3) \otimes U(1)$ spherical harmonics and corresponding Pauli spinors. To our knowledge the construction of these new objects of the Euclidean Taub-NUT geometry has not been well-described in the literature.

In other respects, the four-dimensional Taub-NUT and Taub-NUT-AdS solutions of Einstein's equations play a central role in the construction of diverse and interesting M-theory configurations. The present approach can be easily extended to higher-dimensional spacetimes solutions to the vacuum Einstein equations [31]. There are solutions with non-trivial topology describing spacetimes which are either locally AdS or locally asymptotically flat [32]. Summing up, we believe that the extension of the present approach for the construction of the spherical harmonics to higher-dimensional spacetimes could be relevant within the context of M-theory and deserves further studies.

Acknowledgments

We would like to thank L. Gy. Feher for suggesting to one of us (I.I.C.) a suitable approach to the problem presented here.

Appendix A

The function $h(\beta, \theta, \phi)$

Integrating separately the Eqs. (22) and (23) one obtains [33]

$$h_{(1)}(\beta, \theta, \phi) = F(\beta, \phi) - 2\mu \arctan \left[\frac{\tan \frac{\theta}{2} \tan \frac{\beta}{2} - \cos \phi}{\sin \phi} \right], \quad (\text{A.1})$$

$$h_{(2)}(\beta, \theta, \phi) = \mu\phi + G(\beta, \theta) - 2\mu \arctan \left[\frac{(\cos \theta + \cos \beta) \tan \frac{\phi}{2}}{1 + \cos(\theta + \beta)} \right] \quad (\text{A.2})$$

where F and G are arbitrary functions. If we take $F = 2\mu\phi - \mu\pi$ and $G = 0$ then we find the function

$$\begin{aligned} h(\beta, \theta, \phi) &= h_{(1)}(\beta, \theta, \phi) = h_{(2)}(\beta, \theta, \phi) \\ &= -2\mu \arctan \left[\frac{\sin \phi}{\cot \frac{\theta}{2} \cot \frac{\beta}{2} - \cos \phi} \right] \end{aligned} \quad (\text{A.3})$$

that is just the restriction of the function (28) for $\alpha = \gamma = 0$.

The terms of the angular momentum produced by h have to be calculated according to Eqs. (29) and (8) starting with

$$\left. \frac{\partial}{\partial \beta} h(\beta, \theta, \phi) \right|_{\beta=0} = -\mu \frac{\sin \theta \sin \phi}{1 + \cos \theta}. \quad (\text{A.4})$$

Then, denoting $\xi^2 = \beta$ for $\alpha = \gamma = 0$ and $\xi^3 = \alpha$ for $\beta = \gamma = 0$ and using a simple rotation of angle $\pi/2$ around the third axis one arrives at

$$\left. \frac{\partial h(R, \mathbf{x})}{\partial \xi^{1,2}} \right|_{\xi=0} = -\mu \frac{x^{1,2}}{r + x^3}, \quad \left. \frac{\partial h(R, \mathbf{x})}{\partial \xi^3} \right|_{\xi=0} = -\mu \quad (\text{A.5})$$

where $R = R(\xi)$ is a rotation in the Cayley-Klein parametrization.

References

- [1] S. W. Hawking, *Phys. Lett.* **60A** (1977) 81.
- [2] N. S. Manton, *Phys. Lett.* **B110** (1985) 54; *id*, **B154** (1985) 397; *id*, (E) **B157** (1985) 475.
- [3] M. F. Atiyah and N. Hitchin, *Phys. Lett.* **A107** (1985) 21.
- [4] D. J. Gross and M. J. Perry, *Nucl. Phys.* **B226** (1983) 29.
- [5] R. D. Sorkin, *Phys. Rev. Lett.* **51** (1983) 87.
- [6] G. W. Gibbons and C. A. R. Herdeiro, *Class. Quantum Grav.* **16** (1999) 3619.
- [7] G. W. Gibbons and N. S. Manton, *Nucl. Phys.* **B274** (1986) 183.
- [8] G. W. Gibbons and P. J. Ruback, *Phys. Lett.* **B188** (1987) 226; *Commun. Math. Phys.* **115** (1988) 267.
- [9] L. Gy. Feher and P. A. Horvathy, *Phys. Lett.* **B182** (1987) 183; *id*, (E) **B188** (1987) 512.
- [10] B. Cordani, L. Gy. Feher and P. A. Horvathy, *Phys. Lett.* **B201** (1988) 481.
- [11] K. Yano, *Ann. Math.* **55** (1952) 328.
- [12] J. W. van Holten, *Phys. Lett.* **B342** (1995) 47.
- [13] D. Vaman and M. Visinescu, *Phys. Rev. D* **57** (1998) 3790.
- [14] D. Vaman and M. Visinescu, *Fortschr. Phys* **47** (1999) 493.
- [15] I. I. Cotăescu and M. Visinescu, *Mod. Phys. Lett. A* **15** (2000) 145.
- [16] I. I. Cotăescu and M. Visinescu, *Int. J. Mod. Phys. A* **16** (2001) 1743.
- [17] I. I. Cotăescu and M. Visinescu, *Phys. Lett.* **B502** (2001) 229.
- [18] I. I. Cotăescu and M. Visinescu, *Class. Quantum Grav.* **18** (2001) 3383.
- [19] I. I. Cotăescu and M. Visinescu, *J. Math. Phys.* **43** (2002) 2987.

- [20] G. Mackey, *Induced Representations of Groups and Quantum Mechanics* (Benjamin, New York, 1968).
- [21] W.- K. Tung, *Group Theory in Physics* (World Sci., Philadelphia, 1985).
- [22] B. Carter and R. G. McLenaghan, *Phys. Rev. D* **19** (1979) 1093.
- [23] I. I. Cotăescu, *J. Phys. A: Math. Gen.* **33** (2000) 9177.
- [24] H. Boutaleb - Joutei and A. Chakrabarti, *Phys. Rev. D* **21** (1979) 2280.
- [25] E. Newman and R. Penrose, *J. Math. Phys.* **7** (1966) 863
- [26] T. T. Wu and C. N. Yang, *Nucl. Phys.* **B107** (1976) 365.
- [27] K. Gosh, *Phys. Rev. D* **67** (2003) 124027.
- [28] L. G. Mardoyan, *J. Math. Phys.* **44** (2003) 4981.
- [29] B. Thaller, *The Dirac Equation* (Springer-Verlag, Berlin, Heidelberg, 1992).
- [30] Y. Kazama, C. N. Yang and A. S. Goldhaber, *Phys. Rev.* **15D** (1977) 2287.
- [31] S. Bais and P. Batenberg, *Nucl. Phys.* **B253** (1985) 162.
- [32] A. M. Awad and A. Chamblin, *Clas. Quantum Grav.* **19** (2002) 2051.
- [33] I. S. Gradshteyn and I. M. Ryzhik, *Table of integrals, series and products* (Academic Press, San Diego, 1979).